

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 May 2002 (23.05.2002)

PCT

(10) International Publication Number
WO 02/39958 A2

(51) International Patent Classification: A61K

(21) International Application Number: PCT/US01/51136

(22) International Filing Date:
2 November 2001 (02.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/245,878 3 November 2000 (03.11.2000) US

(71) Applicant (for all designated States except US): TULARIK INC. [US/US]; Two Corporate Drive, South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHWENDNER,

Susan [US/US]; 2000 Crystal Springs Road, #7-05, San Bruno, CA 94066 (US). TIMMERMAN, Pieter [NL/US]; 850 Hillcrest Drive, Redwood City, CA 94062 (US). WALLING, Jacqueline [GB/US]; 2710 Easton Drive, Burlingame, CA 94010 (US).


(74) Agents: KEZER, William, B. et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

[Continued on next page]

(54) Title: COMBINATION THERAPY USING PENTAFLUOROBENZENESULFONAMIDES AND ANTINEOPLASTIC AGENTS

Efficacy of Compound 2 or Gemcitabine Either Alone or in Combination Against MX-1 Human Mammary Tumor Xenografts in Athymic Nude Mice



(57) Abstract: Combination therapies are provided for the treatment of proliferative disorders which use a pentafluorobenzenesulfonamide of formula I and an antineoplastic agent such as gemcitabine or paclitaxel.

WO 02/39958 A2



(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**COMBINATION THERAPY USING
PENTAFLUOROBENZENESULFONAMIDES AND
ANTINEOPLASTIC AGENTS**

5 **CROSS-REFERENCES TO RELATED APPLICATIONS**

This application claims the benefit of USSN 60/245,878, filed November 3, 2000, the disclosure of which is incorporated herein by reference. Also, this application is related in technology to co-pending application Serial Number 09/627,041, filed July 27, 2000.

10

**STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER
FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT**

Not applicable

15

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

20

The present invention relates to combinations of pentafluorobenzenesulfonamides and various other chemotherapeutic agents that are capable of inhibiting abnormal cell proliferation.

BACKGROUND

25

Cancer is a generic name for a wide range of cellular malignancies characterized by unregulated growth, lack of differentiation, and the ability to invade local tissues and metastasize. These neoplastic malignancies affect, with various degrees of prevalence, every tissue and organ in the body. A multitude of therapeutic agents have been developed over the past few decades for the treatment of various types of cancer. The most commonly used types of anticancer agents include: DNA-alkylating agents (e.g., cyclophosphamide, ifosfamide), antimetabolites (e.g., methotrexate, a folate antagonist, and 5-fluorouracil, a pyrimidine antagonist), microtubule disruptors (e.g., vincristine, vinblastine, paclitaxel), DNA intercalators (e.g., doxorubicin, daunomycin),

and hormone therapy (e.g., tamoxifen, flutamide). The ideal antineoplastic drug would kill cancer cells selectively, with a wide therapeutic index relative to its toxicity towards non-malignant cells. It would also retain its efficacy against malignant cells, even after prolonged exposure to the drug. Unfortunately, none of the current chemotherapies
5 possess an ideal profile. Most possess very narrow therapeutic indexes and, in practically every instance, cancerous cells exposed to slightly sublethal concentrations of a chemotherapeutic agent will develop resistance to such an agent, and quite often cross-resistance to several other antineoplastic agents.

The development of new anticancer agents has given rise to new treatment
10 regimens and new combinations that are proving more effective in combating this disease.

Accordingly, it is one object of the present invention to provide compositions which directly or indirectly are toxic to actively dividing cells and are useful in the treatment of cancer.


15 A further object of the present invention is to provide methods for killing actively proliferating cells, such as cancerous, bacterial, or epithelial cells, and treating all types of cancers, and generally proliferative conditions. A further object is to provide methods for treating other medical conditions characterized by the presence of rapidly proliferating cells, such as psoriasis and other skin disorders.

20 Additional objects, features and advantages will become apparent to those skilled in the art from the following description and claims.

SUMMARY OF THE INVENTION

25

In one aspect, the present invention provides compositions useful for the treatment of cancer and other diseases associated with abnormal cell proliferation. The compositions comprise an antineoplastic agent, including but not limited to prodrugs thereof, pharmaceutically acceptable salts of these agents and a compound having the
30 formula:



In the formula above, the letter R represents a hydrogen, substituted or unsubstituted (C₁-C₁₀)alkyl, or substituted or unsubstituted (C₃-C₆)alkenyl. The symbol
 5 Ar represents a substituted or unsubstituted aryl group or a substituted or unsubstituted heteroaryl group.

Suitable antineoplastic or antiproliferative agents include, but are not limited to, DNA-alkylating agents (e.g., cyclophosphamide, BCNU, busulfan and temozolamide), antimetabolites, antifolates and other inhibitors of DNA synthesis (e.g.,
 10 methotrexate, 5-fluorouracil, gemcitabine), microtubule disruptors (e.g., vincristine, vinorelbine, paclitaxel, docetaxel), DNA intercalators (e.g., doxorubicin, daunomycin), hormone agents (e.g., tamoxifen, flutamide), topoisomerase I/II inhibitors and DNA repair agents (e.g., hydroxyurea, camptothecin, etoposide), growth factor receptor kinase inhibitors (e.g., AG1478 and AG1296), biological response modifiers (including
 15 cytokines such as interferon α and growth factor inhibitors), antiangiogenic and antivascular agents (e.g., combretastatin A-4), and other agents such as immunoconjugates (e.g., trasuzamab) and antisense oligonucleotides.

The compositions will, in some embodiments, contain a pharmaceutically acceptable carrier or diluent.

20 In another aspect, the present invention provides methods for the treatment of cancer and other proliferative disorders using the compositions provided above, or using the components in a sequential or simultaneous administration.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 is a graph which illustrates the effects of Compound 2 with gemcitabine in the treatment of MX-1 human mammary tumor xenografts in athymic nude mice, using suboptimal doses of each of the agents.

Figure 2 is a graph which illustrates the effects of Compound 2 with paclitaxel in the treatment of MX-1 human mammary tumor xenografts in athymic nude mice, using suboptimal doses of each of the agents.

Figure 3 provides the structures of Compound 1, Compound 2 and
5 Compound 3.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Definitions

10

The term "alkyl," by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or combination thereof, which may be fully saturated, mono- or polyunsaturated and can include di- and multivalent radicals, having the number of carbon atoms designated (*i.e.*
15 C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)methyl, cyclopropylmethyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl
20 groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. The term "alkyl," unless otherwise noted, is also meant to include those derivatives of alkyl defined in more detail below as "heteroalkyl." Alkyl groups which are limited to hydrocarbon groups are termed "homoalkyl".

25

The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-, and further includes those groups described below as "heteroalkylene." Typically, an alkyl (or
alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower
30 alkylene" is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms.

The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively.

The term "heteroalkyl," by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and from one to three heteroatoms selected from the group consisting of O, N, Si and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S may be placed at any interior position of the heteroalkyl group. The heteroatom Si may be placed at any position of the heteroalkyl group, including the position at which the alkyl group is attached to the remainder of the molecule. Examples include -CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃, -CH₂-CH₂-S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃, -CH₂-CH=N-OCH₃, and -CH=CH-N(CH₃)-CH₃. Up to two heteroatoms may be consecutive, such as, for example, -CH₂-NH-OCH₃ and -CH₂-O-Si(CH₃)₃. Similarly, the term "heteroalkylene" by itself or as part of another substituent means a divalent radical derived from heteroalkyl, as exemplified by -CH₂-CH₂-S-CH₂CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (e.g., alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied.

The terms "cycloalkyl" and "heterocycloalkyl", by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of "alkyl" and "heteroalkyl", respectively. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like.

The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as "haloalkyl," are meant to include monohaloalkyl and polyhaloalkyl. For example, the term "halo(C₁-C₄)alkyl" is meant to include trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

The term "aryl" means, unless otherwise stated, a polyunsaturated, typically aromatic, hydrocarbon substituent which can be a single ring or multiple rings

(up to three rings) which are fused together or linked covalently. The term "heteroaryl" refers to aryl groups (or rings) that contain from zero to four heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

For brevity, the term "aryl" when used in combination with other terms (e.g., aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above. Thus, the term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (e.g., benzyl, phenethyl, pyridylmethyl and the like) including those alkyl groups in which a carbon atom (e.g., a methylene group) has been replaced by, for example, an oxygen atom (e.g., phenoxymethyl, 2-pyridyloxymethyl, 3-(1-naphthyloxy)propyl, and the like).

Each of the above terms (e.g., "alkyl," "heteroalkyl," "aryl" and "heteroaryl") are meant to include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be a variety of groups selected from: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN and -NO₂ in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such radical. R', R'' and R''' each independently refer to hydrogen, unsubstituted (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted alkyl, alkoxy or thioalkoxy groups, or aryl-(C₁-C₄)alkyl groups. When R' and R'' are attached to the same

nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term "alkyl" is meant to include groups such as haloalkyl (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

Similarly, substituents for the aryl and heteroaryl groups are varied and are selected from: -halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R''', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -N₃, -CH(Ph)₂, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'' and R''' are independently selected from hydrogen, (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl.

Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U-, wherein T and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from 0 to 2. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -(CH₂)_s-X-(CH₂)_t-, where s and t are independently integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituent R' in -NR'- and -S(O)₂NR'- is selected from hydrogen or unsubstituted (C₁-C₆)alkyl.

As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient

amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be
5 obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric,
10 hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, oxalic, maleic, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and
15 the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of Pharmaceutical Science*, 1977, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

The neutral forms of the compounds may be regenerated by contacting the
20 salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

In addition to salt forms, the present invention provides compounds which
25 are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the
30 present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of

the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

5 Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are all intended to be encompassed within the scope of the present invention.

10 The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

15

General

20 A number of arylsulfonamides have recently been described for the treatment of disorders and conditions arising from abnormal cell proliferation and from elevated plasma cholesterol levels. See, for example, PCT publications WO 97/30677, WO 98/05315 and WO 99/10320. Representative of this new class of anticancer agents are the pentafluorobenzenesulfonamides described in WO 98/05315. These agents are thought to exert their effect by binding to β -tubulin and disrupting microtubule formation. See, Medina *et al.*, Bioorganic & Med. Chem. Letters, 8(19):2653-56 (1998).

25 Still other pentafluorobenzenesulfonamides have been described in co-pending applications Ser. Nos. 60/090,681 filed June 25, 1998 and 09/336,062 filed June 18, 1999; Ser. Nos. 60/093,570 filed July 20, 1998 and 09/353,976 filed July 15, 1999; and Ser No. 60/100,888 filed September 23, 1998.


30 Clinical trials are in progress to evaluate the pentafluorobenzene-sulfonamide class of compounds for the treatment of cancer, both alone and in combination with other agents. The concept of combination therapy is well exploited in current medical practice. Treatment of a pathology by combining two or more agents that target the same pathogen or biochemical pathway sometimes results in greater efficacy and diminished side effects relative to the use of the therapeutically relevant dose of each

agent alone. In some cases, the efficacy of the drug combination is *additive* (the efficacy of the combination is approximately equal to the sum of the effects of each drug alone), but in other cases the effect can be *synergistic* (the efficacy of the combination is greater than the sum of the effects of each drug given alone). In real medical practice, it is often quite difficult to determine if drug combinations are additive or synergistic.

Description of the Embodiments

Compositions

In one aspect, the present invention provides compositions comprising an antineoplastic agent and a compound having the formula:



(I)

or a pharmaceutically acceptable salt thereof.

In the formula above, the letter R represents a hydrogen, substituted or unsubstituted (C₁-C₁₀)alkyl, or substituted or unsubstituted (C₃-C₆)alkenyl. The symbol Ar represents a substituted or unsubstituted aryl group or a substituted or unsubstituted heteroaryl group.

In preferred embodiments, R represents a hydrogen or a substituted or unsubstituted (C₁-C₄)alkyl group, more preferably hydrogen, methyl or ethyl.

Also preferred are those embodiments in which Ar represents a substituted aryl or substituted heteroaryl group, preferably those having a single ring (e.g., substituted phenyl, substituted pyridyl and substituted pyrimidyl). Particularly preferred embodiments are those in which Ar is substituted phenyl. For those embodiments in which Ar is substituted phenyl, the substituents will typically be present in a number of from one to three. Preferred substituents are selected from -halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR'R''', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, where R', R'' and R''' are independently selected from hydrogen, (C₁-C₄)alkyl, unsubstituted aryl and


heteroaryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl.

Particularly preferred substituents are halogen, (C₁-C₄)alkyl, -OR', -OC(O)R',

-NR'R'', -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R',



-NR'-C(O)NR''R''', perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in which R', R''

- 5 and R''' are hydrogen or (C₁-C₄)alkyl. Still further preferred are those embodiments in which Ar is selected from:




In the most preferred embodiments of the invention, the

- 10 pentafluorobenzenesulfonamide compound used in the composition is selected from:



and



The compositions of the present invention will further comprise an antineoplastic agent. Suitable antineoplastic or antiproliferative agents include, but are not limited to, DNA-alkylating agents (e.g., cyclophosphamide, BCNU, busulfan and temozolamide), antimetabolites, antifolates and other inhibitors of DNA synthesis (e.g., methotrexate, 5-fluorouracil, gemcitabine), microtubule disruptors (e.g., vincristine, vinorelbine, paclitaxel, docetaxel), DNA intercalators (e.g., doxorubicin, daunomycin), hormone agents (e.g., tamoxifen, flutamide), topoisomerase I/II inhibitors and DNA repair agents (e.g., hydroxyurea, camptothecin, etoposide), growth factor receptor kinase inhibitors (e.g., AG1478 and AG1296), biological response modifiers (including cytokines such as interferon α and growth factor inhibitors), antiangiogenic and antivascular agents (e.g., combretastatin A-4), and other agents such as immunoconjugates (e.g., trasuzamab) and antisense oligonucleotides.

Thus, in one embodiment of the present invention, the composition comprises a pentafluorobenzenesulfonamide as defined herein and an antineoplastic agent selected from the group consisting of DNA-alkylating agents, antimetabolites, antifolates and other inhibitors of DNA synthesis, microtubule disruptors, DNA intercalators, hormone agents, topoisomerase I/II inhibitors, DNA repair agents, growth factor receptor kinase inhibitors, biological response modifiers, antiangiogenic and antivascular agents, immunoconjugates and antisense oligonucleotides.

In another embodiment, the composition comprises a pentafluorobenzenesulfonamide as defined herein and an antineoplastic agent selected from the group consisting of cyclophosphamide, BCNU (carmustine), busulfan, temozolomide, UFT, capecitabine, gemcitabine, cytarabine, improsulfan, piposulfan,

benzodepa, carboquone, meturedpa, uredepa, altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate, trimethylolmelamine, chlorambucil, estramustine, ifosfamide, novembrichin, prednimustine, uracil mustard, dacarbazine, fluorouracil, methotrexate, mercaptopurine, thioguanine, vinblastine, vincristine, vinorelbine, vindesine, etoposide, teniposide, daunorubicin, doxorubicin, epirubicin, mitomycin, dactinomycin, daunomycin, plicamycin, bleomycin, L-asparaginase, camptothecin, hydroxyurea, procarbazine, mitotane, aminoglutethimide, tamoxifen, flutamide, mitoxantrone, paclitaxel, docetaxol, and thiotepa.

In preferred embodiments, the antineoplastic agent is gemcitabine or paclitaxel.

As noted above, in the most preferred embodiments of the present invention, the pentafluorobenzenesulfonamide compound used in the compositions is selected from Compound 1, Compound 2, and Compound 3 (see Figure 3). While an understanding of the mechanism by which these compounds are metabolized is not necessary in order to practice the present invention, it is believed that glutathione conjugation plays a major role. Some preferred embodiments of the invention entail the use of compositions comprising Compound 1, Compound 2, or Compound 3 with an antineoplastic agent whose metabolism also is dependent, at least in part, on the formation of a glutathione conjugate (which may include, e.g., BCNU, cyclophosphamide, and thiotepa). Determination of glutathione metabolism can be accomplished according to standard methods known to those of skill in the art (see, e.g., Mannervik and Widersten in ADV. IN DRUG METAB. IN MAN, G.M. Pacifici and G.N. Fracchis, eds., European Commission, Luxemburg: 407-459 (1995), using glutathione transferases available from commercial sources such as PanVera, product nos. P2175, P2192 and P2177, and Research Diagnostics). Enhanced efficacy may be observed with such combinations due to competition for glutathione metabolism. Depending on the agents involved, this may result in depletion of glutathione levels, delayed metabolism of one or both agents, and increased exposure of the malignant tissue to one or more of the composition's active components.

Methods of Treating Proliferative Disorders

The present invention provides, in another aspect, methods for the treatment of proliferative disorders. In one embodiment, treatment is carried out using a

composition comprising each of the two agents described above. In another embodiment, treatment comprises separate administration of one or more antineoplastic agents and a pentafluorophenylsulfonamide of formula I.

5 i. combination composition

In this embodiment of the invention, a composition of two or more agents (described above) is administered to a patient in need of treatment. The amount of each agent will typically be less than an amount that would produce a therapeutic effect if administered alone. The precise method of administration will depend on the patient and
10 the judgment of the clinician, but will preferably be intravenous.

 ii. compositions used sequentially (administer each separately)

In this embodiment of the invention, conventional protocols are described for the administration of an antineoplastic agent and compound 1 (as representative of the compounds of formula I). One of skill in the art will understand that various changes can
15 be made by the clinician, depending on the particular agents selected for use and the routes and timing of administration. Thus, the present invention contemplates that the antineoplastic agent and the compounds of formula I can be administered sequentially on the same day, on concurrent days, or up to about 4 weeks apart.

The antineoplastic agent is preferably administered with a single
20 intravenous infusion on day one of compound 1 administration period about four hours after the first day's administration of compound 1. To maintain sufficient hydration, one liter of normal saline with 20 meq KCl/L and 1 gm of magnesium sulfate, at a rate of about 250 ml/hour is administered prior to and after the infusion. Additional fluid may be given to maintain adequate urine output.

25 The treatment cycle may be continued until a clinical response is achieved or until intolerable side effects are encountered. The dosages of compound 1 and/or antineoplastic agent may be increased with each new treatment cycle, provided intolerable side effects are not encountered. The dosages may also be decreased if intolerable side effects are encountered. It is presently preferred to gradually adjust the
30 dosage of compound 1 while holding the antineoplastic agent dosage constant.

As alluded to previously, certain preferred embodiments of the present invention entail combination therapy involving a pentafluorobenzenesulfonamide compound selected from Compound 1, Compound 2, and Compound 3 (see Figure 3) and at least one other antineoplastic agent, wherein metabolism of the other antineoplastic

agent(s) is dependent, at least in part, on the formation of a glutathione conjugate. In such embodiments, the order of administration may be especially important; that is, the order of administration may result in enhanced efficacy while minimizing adverse effects. In some preferred embodiments, it is preferable to administer Compound 1, Compound 2
5 or Compound 3 prior to the other antineoplastic agent, while in other preferred embodiments it is advantageous to co-administer the agents.

A common, but tolerable side effect of antineoplastic agent is nausea and vomiting. This can be alleviated by administering an anti-emetic (e.g., Ondansetron®, Granisetron®, Decadron®, Haldol®, Benadryl®, Ativan® and the like).

10 Of course, other forms of administration of both active ingredients, as they become available, are contemplated, such as by nasal spray, transdermally, by suppository, by sustained release dosage form, by IV injection, etc. Any form of administration will work so long as the proper dosages are delivered without destroying the active ingredient.

15 The effectiveness of treatment may be determined by controlled clinical trials. Patients having cancer with measurable or evaluable tumors will be included in the study. A measurable tumor is one that can be measured in at least two dimensions such as a lung tumor surrounded by aerated lung, a skin nodule, or a superficial lymph node. An evaluable tumor is one that can be measured in one dimension such as a lung tumor
20 not completely surrounded by aerated lung or a palpable abdominal or soft tissue mass that can be measured in one dimension. Tumor markers which have been shown to be highly correlated with extent of disease will also be considered to provide an evaluable disease, such as PSA for prostate cancer, CA-125 for ovarian cancer, CA-15-3 for breast cancer, etc.

25 The tumor will be measured or evaluated before and after treatment by whatever means provides the most accurate measurement, such as CT scan, MRI scan, Ultrasonography, etc. New tumors or the lack thereof in previously irradiated fields can also be used to assess the anti-tumor response. The criteria for evaluating response will be similar to that of the WHO Handbook of Reporting Results of Cancer Treatment,
30 WHO Offset Publication 1979, 49-World Health Organization, Geneva. The following results are defined for uni- and bi-dimensionally measurable tumors.

Complete response: Complete disappearance of all clinically detectable malignant disease determined by two observations not less than four weeks apart.

Partial Response: (a) for bidimensionally measurable tumors, a decrease of

at least 50% in the sum of the products of the largest perpendicular diameters of all measurable tumors as determined by two observations not less than four weeks apart. (b) for unidimensionally measurable tumors, a decrease by at least 50% in the sum of the largest diameters of all tumors as determined by two observations not less than four weeks apart. In cases where the patient has multiple tumors, It is not necessary for all tumors to have regressed to achieve a partial response as defined herein, but no tumor should have progressed and no new tumor should appear.

Stable disease: (a) for bidimensionally measurable tumors, less than a 50% decrease to less than a 25% increase in the sum of the products of the largest perpendicular diameters of all measurable tumors. (b) for unidimensionally measurable tumors, less than a 50% decrease to less than a 25 % increase in the sum of the diameters of all tumors. For (a) and (b) no new tumors should appear.

No clinical response, i.e. progressive disease is defined as an increase of more than 50% in the product of the largest perpendicular diameters for at least one bidimensionally measurable tumor, or an increase of more than 25% in measurable dimension of at least one unidimensionally measurable tumor.

Of course elimination or alleviation of other known signs or symptoms of cancer, especially those listed previously can also be used to evaluate the effectiveness of this invention.

The cancers should be evaluated, i.e. tumors measured, etc., no more than 14 days before the start of the treatment. These cancers should be reevaluated about 28 days after day 1 of administration of the first dose of compound 1 and antineoplastic agent. Twenty eight days after this initial administration another administration period may be performed, and evaluations performed 28 days after the start of this second cycle. The treatment cycles may be continued until a clinical response is achieved or unacceptable toxicity is encountered.

Another aspect of this invention is the treatment of cancer with reduced side effects normally associated with an antineoplastic agent. This objective can be achieved by administration of lower doses of the two active ingredients or by shorter duration of dosing brought about by the synergistic effect of the combination.

EXAMPLES


Figures 1 and 2 illustrate the effect achieved by combining a pentafluorobenzenesulfonamide with gemcitabine or with paclitaxel.

5

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example
10 for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

WHAT IS CLAIMED IS:

1 1. A composition for the treatment of proliferative disorders,
2 comprising an antineoplastic agent and a compound having the formula:



3
4 and pharmaceutically acceptable salts thereof;
5 wherein

6 R is a member selected from the group consisting of hydrogen and
7 substituted or unsubstituted (C₁-C₁₀)alkyl; and

8 Ar is a member selected from the group consisting of substituted or
9 unsubstituted aryl and substituted or unsubstituted heteroary

1 2. A composition in accordance with claim 1, wherein said
2 antineoplastic agent is selected from the group consisting of DNA-alkylating agents,
3 antimetabolites, antifolates and other inhibitors of DNA synthesis, microtubule disruptors,
4 DNA intercalators, hormone agents, topoisomerase I/II inhibitors, DNA repair agents,
5 growth factor receptor kinase inhibitors, biological response modifiers, antiangiogenic
6 and antivasular agents, immunoconjugates and antisense oligonucleotides.

1 3. A composition in accordance with claim 1, wherein said
2 antineoplastic agent is selected from the group consisting of cyclophosphamide, BCNU,
3 busulfan, temozolomide, UFT, capecitabine, gemcitabine, cytarabine, improsulfan,
4 piposulfan, benzodepa, carboquone, meturedpa, uredepa, altretamine,
5 triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate,
6 trimethylolmelamine, chlorambucil, estramustine, ifosfamide, novembrichin,
7 prednimustine, uracil mustard, dacarbazine, fluorouracil, methotrexate, mercaptopurine,
8 thioguanine, vinblastine, vincristine, vinorelbine, vindesine, etoposide, teniposide,
9 daunorubicin, doxorubicin, epirubicin, mitomycin, dactinomycin, daunomycin,
10 plicamycin, bleomycin, L-asparaginase, camptothecin, hydroxyurea, procarbazine,
11 mitotane, aminoglutethimide, tamoxifen, flutamide, mitoxantrone, paclitaxel, docetaxol,
12 and thiotepa.

13 4. A composition in accordance with claim 1, wherein said
14 antineoplastic agent is selected from the group consisting of doxorubicin, daunorubicin,
15 gemcitabine and paclitaxel.


16 5. A composition in accordance with claim 1, wherein said
17 antineoplastic agent is gemcitabine or paclitaxel.

1 6. A composition in accordance with claim 1, wherein R is hydrogen
2 or unsubstituted (C₁-C₄)alkyl.


1 7. A composition in accordance with claim 1, wherein Ar is a
2 substituted phenyl group.

1 8. A composition in accordance with claim 7, wherein said
2 substituents on said phenyl group are selected from the group consisting of halogen, (C₁-
3 C₄)alkoxy, (C₁-C₄)alkyl, -OPO₃H₂,

1 9. A composition in accordance with claim 8, wherein Ar represents a
2 member selected from the group consisting of




- 1 10. A composition in accordance with claim 1, wherein said compound
2 is selected from the group consisting of:



1 11. A method for the treatment of a proliferative disorder, comprising
 2 administering to a subject in need of such treatment an effective amount of a composition
 3 of claim 1.

1 12. A. method in accordance with claim 11, wherein said compound is
 2 selected from the group consisting of:



4
 5
 6 13. A method in accordance with claim 12, wherein said antineoplastic
 7 agent is selected from the group consisting of DNA-alkylating agents, antimetabolites,
 8 antifolates and other inhibitors of DNA synthesis, microtubule disruptors, DNA
 9 intercalators, hormone agents, topoisomerase I/II inhibitors, DNA repair agents, growth
 10 factor receptor kinase inhibitors, biological response modifiers, antiangiogenic and
 11 antivascular agents, immunoconjugates and antisense oligonucleotides.

1 14. A method in accordance with claim 12, wherein said antineoplastic
 2 agent is selected from the group consisting of cyclophosphamide, BCNU, busulfan,
 3 temozolomide, UFT, capecitabine, gemcitabine, cytarabine, improsulfan, pipsulfan,
 4 benzodepa, carboquone, meturedpa, uredepa, altretamine, triethylenemelamine,
 5 triethylenephosphoramidate, triethylenethiophosphoramidate, trimethylolmelamine,
 6 chlorambucil, estramustine, ifosfamide, novembrichin, prednimustine, uracil mustard,
 7 dacarbazine, fluorouracil, methotrexate, mercaptopurine, thioguanine, vinblastine,

8 vincristine, vinorelbine, vindesine, etoposide, teniposide, daunorubicin, doxorubicin,
 9 epirubicin, mitomycin, dactinomycin, daunomycin, plicamycin, bleomycin, L-
 10 asparaginase, camptothecin, hydroxyurea, procarbazine, mitotane, aminoglutethimide,
 11 tamoxifen, flutamide, mitoxantrone, paclitaxel, docetaxol, and thiotepa.


12 15. A method in accordance with claim 12, wherein said antineoplastic
 13 agent is selected from the group consisting of doxorubicin, daunorubicin, gemcitabine
 14 and paclitaxel.

15 16. A method in accordance with claim 12, wherein said antineoplastic
 16 agent is gemcitabine or paclitaxel.

17 17. A method for the treatment of a proliferative disorder, comprising
 18 administering to a subject in need of such treatment:

19 i) a first amount of an antineoplastic agent; and

20 ii) a second amount of a compound of formula:





21 and pharmaceutically acceptable salts thereof; wherein

22 R is a member selected from the group consisting of hydrogen and
 23 substituted or unsubstituted (C₁-C₁₀)alkyl; and


24 Ar is a member selected from the group consisting of substituted or
 25 unsubstituted aryl and substituted or unsubstituted heteroaryl;

26 wherein said first amount and said second amount, in combination, are
 27 effective to treat said proliferative disorder

28 18. A method in accordance with claim 17, wherein said compound is
 1 selected from the group consisting of
 2



and



3

4

5 19. A method in accordance with claim 18, wherein said antineoplastic
6 agent is selected from the group consisting of DNA-alkylating agents, antimetabolites,
7 antifolates and other inhibitors of DNA synthesis, microtubule disruptors, DNA
8 intercalators, hormone agents, topoisomerase I/II inhibitors, DNA repair agents, growth
9 factor receptor kinase inhibitors, biological response modifiers, antiangiogenic and
10 antivascular agents, immunoconjugates and antisense oligonucleotides.

1 20. A method in accordance with claim 18, wherein said antineoplastic
2 agent is selected from the group consisting of cyclophosphamide, BCNU, busulfan,
3 temozolomide, UFT, capecitabine, gemcitabine, cytarabine, improsulfan, pipsulfan,
4 benzodepa, carboquone, meturedpa, uredepa, altretamine, triethylenemelamine,
5 triethylenephosphoramidate, triethylenethiophosphoramidate, trimethylolmelamine,
6 chlorambucil, estramustine, ifosfamide, novembrichin, prednimustine, uracil mustard,
7 dacarbazine, fluorouracil, methotrexate, mercaptopurine, thioguanine, vinblastine,
8 vincristine, vinorelbine, vindesine, etoposide, teniposide, daunorubicin, doxorubicin,
9 epirubicin, mitomycin, dactinomycin, daunomycin, plicamycin, bleomycin, L-
10 asparaginase, camptothecin, hydroxyurea, procarbazine, mitotane, aminoglutethimide,
11 tamoxifen, flutamide, mitoxantrone, paclitaxel, docetaxol, and thiotepa.

12 21. A method in accordance with claim 18, wherein said antineoplastic
13 agent is selected from the group consisting of doxorubicin, daunorubicin, gemcitabine
14 and paclitaxel.

15 22. A method in accordance with claim 18, wherein said antineoplastic
16 agent is gemcitabine or paclitaxel.

17

18 23. A method in accordance with claim 18, wherein said antineoplastic
19 agent is administered prior to said compound.

20


21 24. A method in accordance with claim 18, wherein said antineoplastic
22 agent is administered after said compound.

23

24 25. A method in accordance with claim 18, wherein said antineoplastic
25 agent is administered simultaneously with said compound.

FIGURE 1

Efficacy of Compound 2 or Gemcitabine Either Alone or in Combination Against MX-1 Human Mammary Tumor Xenografts in Athymic Nude Mice



Arrow indicates dose administration. Results are expressed as the mean \pm SEM.

FIGURE 2

Efficacy of Compound 2 or Paclitaxel Either Alone or in Combination Against MX-1 Human Mammary Tumor Xenografts in Athymic Nude Mice







FIGURE 3

COMPOUND 1
(sodium salt)

COMPOUND 2



COMPOUND 3

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 May 2002 (23.05.2002)

PCT

(10) International Publication Number
WO 02/039958 A3

(51) International Patent Classification⁷: A61K 31/415,
A01N 43/38, C07D 231/56, C07C 303/00, 307/00

(21) International Application Number: PCT/US01/51136

(22) International Filing Date:
2 November 2001 (02.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/245,878 3 November 2000 (03.11.2000) US

(71) Applicant (for all designated States except US): TU-
LARIK INC. [US/US]; Two Corporate Drive, South San
Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHWENDNER,
Susan [US/US]; 2000 Crystal Springs Road, #7-05,
San Bruno, CA 94066 (US). TIMMERMAN, Pieter
[NL/US]; 850 Hillcrest Drive, Redwood City, CA 94062
(US). WALLING, Jacqueline [GB/US]; 2710 Easton
Drive, Burlingame, CA 94010 (US).


(74) Agents: KEZER, William, B. et al.; Townsend and
Townsend and Crew LLP, Two Embarcadero Center,
Eighth Floor, San Francisco, CA 94111 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,

[Continued on next page]

(54) Title: COMBINATION THERAPY USING PENTAFLUOROBENZENESULFONAMIDES AND ANTINEOPLASTIC AGENTS

**Efficacy of Compound 2 or Gemcitabine Either Alone
or in Combination Against MX-1 Human Mammary
Tumor Xenografts in Athymic Nude Mice**



Arrow indicates dose administration. Results are expressed as the mean \pm SEM.

(57) Abstract: Combination therapies are provided for the treatment of proliferative disorders which use a pentafluorobenzenesulfonamide of formula I and an antineoplastic agent such as gemcitabine or paclitaxel.

WO 02/039958 A3



SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.

Published:

— with international search report

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) **Date of publication of the international search report:**
6 September 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/51136

| | | |
|---|--|--|
| A. CLASSIFICATION OF SUBJECT MATTER | | |
| IPC(7) : A61K 31/415; A01N 43/38; C07D 231/56; C07C 303/00; C07C 307/00 | | |
| US CL : 514/403, 415, 518, 602, 604; 548/361.1, 469; 558/56, 61; 564/90, 92 | | |
| According to International Patent Classification (IPC) or to both national classification and IPC | | |
| B. FIELDS SEARCHED | | |
| Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/403, 415, 518, 602, 604; 548/361.1, 469; 558/56, 61; 564/90, 92 | | |
| Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched | | |
| Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet | | |
| C. DOCUMENTS CONSIDERED TO BE RELEVANT | | |
| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | WO 98/05315 A1 (TULARIK, INC.) 12 February 1998 (12.2.1998), see entire document, especially page 29, lines 21 and 22, and Examples 6 and 7 on page 33. | 1-25 |
| Y | US 5,880,151 A (MEDINA et al) 09 March 1999 (9.3.1999), see entire document, especially column 23, lines 25-55. | 1-25 |
| Y | US 6,121,304 A (FLYGARE et al) 19 September 2000 (19.9.2000), see entire document, especially column 24, lines 1-35. | 1-25 |
| Y | MEDINA et al. Novel antineoplastic agents with efficacy against multidrug resistant tumor cell lines. Bioorg. Med. Chem. Lett. 1998. Vol. 8, No. 9, pages 2653-2656, see entire document, especially abstract. | 1-25 |
| <input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex. | | |
| * Special categories of cited documents: | | |
| "A" | document defining the general state of the art which is not considered to be of particular relevance | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "E" | earlier application or patent published on or after the international filing date | "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "L" | document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "O" | document referring to an oral disclosure, use, exhibition or other means | "&" document member of the same patent family |
| "P" | document published prior to the international filing date but later than the priority date claimed | |
| Date of the actual completion of the international search 18 March 2002 (18.03.2002) | | Date of mailing of the international search report 20 MAY 2002 |
| Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230 | | Authorized officer Richard Schnizer Telephone No. (703) 308-0196 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/51136

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest: ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/51136

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-10, drawn to compositions for the treatment of proliferative disorders.

Group II, claim(s) 11-25, drawn to methods of treating proliferative disorders.

The inventions listed as Groups I and II do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The technical feature linking the two groups cannot be a special technical feature under PCT Rule 13.2 because the technical feature fails to make a contribution over the prior art.

The technical feature linking groups I and II is a pentafluorosulfonamide compound in which the sulfonamide nitrogen is substituted with an aryl group and either a hydrogen or an alkyl group. However, such compounds were known in the prior art. See e.g. WO 98/05315, US Patents 5,880,151, and 6,121,304, and Medina et al (Bioorg. Med. Chem. Lett. (1988) Vol. 8, No. 19, pages 2653-2656.

Because the technical feature linking the inventions was known in the prior art, it cannot be a special technical feature under PCT Rule 13.2, and the claimed inventions lack unity under PCT Rule 13.1.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

The species are as follows: the antineoplastic agents listed in claims 2, 3, 14, and 20.

The species listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The combination of an antineoplastic agent and the pentafluorosulfonamide does not relate to a single general inventive concept under PCT Rule 13.1 because it fails to make a contribution over the prior art. WO 98/05315 teaches this combination. See e.g. page 29, lines 21 and 22; and Examples 6 and 7 on page 33.

However, an election will not be required as the species are considered to be obvious in view of each other.

Continuation of B. **FIELDS SEARCHED** Item 3:
STN: BIOSIS CAPLUS EMBASE BIOTECHDS MEDLIN
WEST